

THE DOMINANT WEEDS IN INTENSIVE ORCHARDS

Branka Ljevnaić-Mašić, Slavko Gudurić, Ljiljana Nikolić, Dejana Džigurski, Aleksandra Petrović

*Faculty of Agriculture, University of Novi Sad, Trg D. Obradovića 8, 21000 Novi Sad, Serbia
e-mail: brana@polj.uns.ac.rs*

Abstract

The main prerequisite for the proper and timely implementation of measures of weed control and their suppression is the knowledge of their basic biological characteristics. Therefore, the aim of this study was floristic and taxonomic analysis, and also analysis of the biological characteristics of the dominant weeds in intensive orchards, which are cultivated according to the principles of conventional agricultural production, around the Irig village (Vojvodina Province, Serbia). It was identified a total of 18 weed species. Classes Magnoliopsida and Liliopsida are preset with 83.33% and 16.67% species, respectively. In the analyzed weed flora were recorded the presence of 16 invasive species for Europe and 4 invasive species for the Vojvodina Province. The largest number of species are weed-ruderal plants (50.00%), followed by ruderal weeds (22.22%). Majority of the analyzed weed species bloomed from June to September. The biological spectrum of the flora indicates the dominance of therophytes (T - 61.10%).

Introduction

Natural conditions in Serbia are extremely favorable for growing fruit [1]. Serbia is one of the few countries, that at a relatively small area, has favorable environmental conditions for the cultivation of all continental species of fruit trees [1, 2]. Fruits are a food of high nutritional value and significant raw material for different products. Fruit growing also allows a rational use of agricultural land. A larger area under the fruit trees positively influences to the microclimate of the place and prevents soil erosion on slopes.

However, one of the biggest problems in agricultural production is a control of weeds. The weeds in orchards inflict great damage during production. Their presence significantly reduces the yield and impairs the quality of the fruit. In intensive production, the aim is to destroy weeds or at least brought them into the balance with the crop [3]. The composition of weeds in orchards is varied making it difficult to correct choice of herbicides for their control. Great influence on the efficiency of most herbicides have climatic factors. Better results are obtained under conditions of higher soil moisture and pre-treated soil [4]. The basic prerequisite for the choice of appropriate measures of weed control is the knowledge of the biological and ecological characteristics of weeds [5, 6, 7, 8, 9]. Therefore, the aim of this study was floristic and taxonomic analysis, and also analysis of basic biological characteristics of the dominant weed species in the orchards around the Irig village (Vojvodina Province, Serbia), cultivated according to the principles of conventional agriculture.

Experimental

Investigation of weed flora in intensive orchards of plums, peaches and apricots, were carried out during the vegetation period 2013-2014, around the Irig village ((Vojvodina Province, Serbia). Weed species were determined according to standard identification keys [10, 11, 12, 13]. The taxonomic affiliation of species is according to [14]. Characterization of invasive species for Europe and for Vojvodina Province are according to [15] and [16]. The categorization of weeds by habitat type is according to [17], and for a species *Consolida*

regalis Gray according to [18]. Time of flowering is according to [17], and for a species *Consolida regalis* Gray according to [19]. Life forms are according to [20].

Investigation area

Irig village (45°10' 19°86') is located in the north of the Republic of Serbia, in terms of moderate continental climate. Summers are very hot, with a lack of moisture, and winters are long and harsh. Autumn and spring are moderately warm and short [21]. The average air temperature is 10.9 °C.

Results and discussion

In the weed flora of intensive orchards was identified a total of 18 species. All weeds are included into the phylum Magnoliophyta (11 families, 10 orders, 8 superorders, 7 subclasses and 2 classes), Tab. 1. The greatest number of weeds belongs to the class Magnoliopsida (83.33%). The most common are the representatives of the family Asteraceae (27.78%) and Poaceae (16.67%). In the analyzed weed flora were recorded the presence of 16 invasive species for the Europe (88.88%) according to [15]. Four species (22.22%) are invasive for the Vojvodina Province [16]: *Amaranthus retroflexus* L., *Consolida regalis* Gray, *Panicum crus-galli* L. and *Stenactis annua* L. (Tab. 1). Considering on the huge problems that invasive weed species do to grown plants, it is necessary constantly monitoring of their numbers which is very important in order to prevent their uncontrolled proliferation. Special attention must be paid to adventitious and cosmopolitan weeds because their intensive spread makes a big problems on grown plants and native species in general [22]. The largest number of weed species belongs to weed-ruderal plants (50.00%), followed by ruderal (22.22%) and segetal weed species (16.67%). The least number of species belong to meadow weeds (11.11%), Tab. 1. Majority of the analyzed weed species bloomed from June (VI) to September (IX), Tab. 1. The longest flowering period has *Capsella bursa-pastoris* (L.) Medik. (IV-XI). The shortest flowering period have *Rubus caesius* L., *Panicum crus-galli* L. and *Calamagrostis epigeios* (L.) Roth. The biological spectrum indicates on the therophytic character of the weed flora (therophytes - 61.10%), Tab. 1. The most abundant are therophytes T₄, i.e. annuals plants which germinate in the Spring, with the maturing at the end of Summer (33.33%). Geophytes (22.22%) and hemicryptophytes (16.67%) are also recorded. Similar results were obtained by [3]. The domination of therophytes indicates on instability of weed community due to the intensive agro-technical measures in the investigated agroecosystem [23, 5].

Conclusion

Investigation of weed flora in intensive orchards around Irig village, cultivated according to the principles of conventional agriculture, it was found a total of 18 weed species. Sixteen species are invasive for Europe and four for Vojvodina Province. All species are included into phylum Magnoliophyta (class Magnoliopsida - 83.33% and class Liliopsida - 16.67%). The most common are the representatives of the family Asteraceae (27.78%). Majority of the analyzed weed species belongs to weed-ruderal plants (50.00%). The greatest number of recorded weed species bloomed from June to September. The study weed flora shows the therophytic character (T - 61.10%) with prevalence of annual plants that germinate in the Spring, with the maturing at the end of Summer (T₄ - 33.33%).

Acknowledgements

This study is part of the project TR31027 titled »Organic agriculture: Improvement of production by use of fertilizers, biopreparates and biological measures» subsidized by the Ministry for Education and Science of the Republic of Serbia.

Table 1. The dominant weed species in intensive orchards with categorization according to the habitat, time of flowering and life forms (Irig village, Serbia)

Phylum	Class	Subclass	Superorder	Order	Familia	Plant species	Categorization according to the habitat	Time of flowering	Life form
Magnoliophyta	Magnoliopsida	Ranunculidae	Ranunculanae	Ranunculales	Ranunculaceae	<i>Consolida regalis</i> Gray *	S	V-IX	T ₂
				Papaverales	Papaveraceae	<i>Papaver rhoeas</i> L. *	S	V-VI	T ₂
		Caryophyllidae	Caryophyllanae	Caryophyllales	Amaranthaceae	<i>Amaranthus retroflexus</i> L. *	KR	VI-IX	T ₄
					Chenopodiaceae	<i>Chenopodium album</i> L. *	KR	VI-XI	T ₄
			Polygonanae	Polygonales	Polygonaceae	<i>Polygonum aviculare</i> L. *	R, KR	V-X	T ₄
		Rosidae	Rosanae	Rosales	Rosaceae	<i>Rubus caesius</i> L.	KR	V	H ₃
		Asteridae	Asteranae	Asterales	Asteraceae	<i>Artemisia vulgaris</i> L. *	R, LK, ŠK	VII-IX	H ₅
						<i>Erigeron canadensis</i> L. *	R	VI-X	T ₄
						<i>Cirsium arvense</i> (L.) Scop. *	KR	VI-VII	G ₃
						<i>Stenactis annua</i> L. *	R	VI-X	T ₄
						<i>Taraxacum officinale</i> L. *	KR, LK	IV-IX	H ₃
		Dilleniidae	Capparanae	Capparales	Brassicaceae	<i>Capsella bursa-pastoris</i> (L.) Medik. *	KR	IV-XI	T ₁
						<i>Sinapis arvensis</i> L. *	KR	V-IX	T ₃
		Lamiidae	Lamianae	Solanales	Convolvulaceae	<i>Convolvulus arvensis</i> L. *	KR	VI-IX	G ₃
				Lamiales	Lamiaceae	<i>Mentha arvensis</i> L. *	LK, ŠK	VI-IX	G ₂
	Liliopsida	Commelinidae	Poanae	Poales	Poaceae	<i>Bromus commutatus</i> Schrad. *	KR	V-VI	T ₂
						<i>Calamagrostis epigeios</i> (L.) Roth.	LK, ŠK	VII	G ₁
						<i>Panicum crus-galli</i> L. *	S	VI	T ₄
Σ	2	7	8	10	11	18			

Legend: *- invasive species for the Europe; bold - invasive species for the Vojvodina Province; Legend: KR - weed-ruderal plant; R – ruderal weed; LK – meadow weed; ŠK – forest weed; S – segetal weed; I-XII – month; T – therophyte; G – geophyte; H - hemicryptophyte

References

- [1] D. Nikolić, Z. Keserović, N. Magazin, S. Paunović, R. Miletić, M. Nikolić, J. Milivojević. Stanje i perspektive razvoja voćarstva u Srbiji. Zbornik radova i apstrakata XIV Kongresa voćara i vinogradara Srbije sa međunarodnim učešćem. Vrnjačka Banja, Srbija. (2012), 3-22.
- [2] Statistički godišnjak Republike Srbije. (2015)(available at: <http://webrzs.stat.gov.rs/WebSite/userFiles/file/Aktuelnosti/StatGod2015.pdf>)
- [3] D. Džigurski, B. Ljevnaić-Mašić, Lj. Nikolić. Dominantni korovi pri organskoj proizvodnji breskve - *Prunus persica* Batsch. (Rosaceae A. L. De Jussieu 1789, Rosales). Zbornik izvoda XXVI Nacionalne konferencije Procesna tehnika i energetika u poljoprivredi - PTEP 2014, 6-11. april 2014, Kladovo, Srbija. (2014), 39-40.
- [4] J. Jocković J. Suzbijanje korova u voćnjacima. Bilten Poljoprivredne stručne službe, Šabac. (2010), 1-5.
- [5] D. Kovačević. Njivski korovi-biologija i suzbijanje. Poljoprivredni fakultet, Univerzitet u Beogradu. (2008), 1-506.
- [6] Lj. Nikolić, D. Džigurski, B. Ljevnaić-Mašić, R. Čabilovski, M. Manojlović. Bulgarian Journal of Agricultural Science. (2011), 17(6), 736-743.
- [7] B. Ljevnaić-Mašić, Lj. Nikolić, D. Džigurski. Acta Biologica Jugoslavica, serija G: Acta herbologica. (2012), 21(1), 41-50.
- [8] D. Džigurski D., Lj. Nikolić, B. Ljevnaić-Mašić. Journal on Processing and Energy in Agriculture. (2013), 17(3), 130-133.
- [9] B. Ljevnaić-Mašić, D. Džigurski, Lj. Nikolić. Journal on Processing and Energy in Agriculture. (2013), 17(1), 33-38.
- [10] M. Josifović (ed.). Flora SR Srbije. I-IX, SANU, Beograd. (1970-1977)
- [11] M. Sarić (ed.). Flora Srbije. X, SANU, Beograd. (1986)
- [12] M. Sarić (ed.). Flora Srbije I. SANU, Beograd. (1992)
- [13] S. Jávorka and V. Csapody Icanographie der Flora des Südostlichen Mitteleuropa. Akademiai Kiado, Budapest. (1975)
- [14] A. Takhtajan. Flowering Plants. Second Edition. Springer. (2009), 1-906.
- [15] DAISIE. Delivering Alien Invasive Species Inventories for Europe.(2016) (available at: <http://www.europe-aliens.org/speciesSearch.do>)
- [16] IASV.Lista invazivnih vrsta na području AP Vojvodine = List of invasive species in AP Vojvodina [Internet]. Version 0.1 beta. Anačkov G, Bjelić-Čabrilo O, Karaman I, Karaman M, Radenković S, Radulović S, Vukov D & Boža P, editors. Department of Biology and Ecology. Novi Sad, Serbia. (2011) (available at: <http://iasv.dbe.pmf.uns.ac.rs>)
- [17] M. Čanak, S. Parabućski, M. Kojić. Ilustrovana korovska flora Jugoslavije. Matica srpska, Novi Sad, Srbija. (1978)
- [18] M. Kojić, A. Stanković, M. Čanak. Korovi - biologija i suzbijanje, Institut za zaštitu bilja, Poljoprivredni fakultet, Novi Sad, Srbija. (1972)
- [19] J. Kovačević. Korovi u poljoprivredi - Herbicidi. Nakladni zavod Znanje. Zagreb. (1976), 1-750.
- [20] M. Ujvárosi. Gymnövények. Mezőgazdasági Kiado, Budapest, Hungary. (1973)
- [21] P. Katić, D. Đukanović, P. Đaković. Klima SAP Vojvodine. Poljoprivredni fakultet, Novi Sad, OOUR Institut za ratarstvo i povrtarstvo, Novi Sad. (1979), 1-237.
- [22] S. Vrbničanin, B. Karadžić, Z. Dajić-Stevanović. Adventivne i invazivne korovske vrste na području Srbije. Acta Herbologica. (2004), 13(1), 1-12.
- [23] M. Kojić, B. Šinžar, R. Stepić. Korovska vegetacija severozapadne Srbije. Zbornik radova Poljoprivrednog fakulteta. (1988), 32-33(589), 83-107.